
EFDA–JET–PR(04)04

V.G. Kiptily, Yu.F. Baranov, R. Barnsley, L. Bertalot, N.C. Hawkes,
A. Murari, S. Popovichev, S.E. Sharapov, D. Stork, V. Yavorskij
and JET EFDA contributors

First Gamma-Ray Measurements of Fusion Alpha Particles in JET Trace Tritium Experiments

First Gamma-Ray Measurements of Fusion Alpha Particles in JET Trace Tritium Experiments

V.G. Kiptily¹, Yu.F. Baranov¹, R. Barnsley¹, L. Bertalot², N.C. Hawkes¹,
A. Murari³, S. Popovichev¹, S.E. Sharapov¹, D. Stork¹, V. Yavorskij^{4,5}
and JET EFDA contributors*

¹*Euratom/UKAEA Fusion Association, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK*

²*Associazione Euratom/ENEA/CNR sulla Fusione, Frascati, Rome, Italy*

³*Consorzio RFX - Associazione Euratom-Enea sulla Fusione, I-35127 Padova, Italy*

⁴*Euratom/OEAW Association, Institute for Theoretical Physics, University of Innsbruck, Austria*

⁵*Institute for Nuclear Research, Kiev, Ukraine*

* See annex of J. Pamela et al, "Overview of Recent JET Results and Future Perspectives",
Fusion Energy 2002 (Proc. 19th Fusion Energy Conference, Lyon, 2002).

“This document is intended for publication in the open literature. It is made available on the understanding that it may not be further circulated and extracts or references may not be published prior to publication of the original when applicable, or without the consent of the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

“Enquiries about Copyright and reproduction should be addressed to the Publications Officer, EFDA, Culham Science Centre, Abingdon, Oxon, OX14 3DB, UK.”

ABSTRACT.

Gamma-ray spectra from nuclear reactions between fusion-born alpha (α) particles and Be impurities were measured for the first time in deuterium plasmas in the Joint European Torus. The time dependence of the measured spectra allowed the determination of the density evolution of fast α particles. Correlation between the decay time of the γ -ray emission and the plasma parameters in different plasma scenarios was established. Results of the measurements are consistent with classical slowing down of the α particles in discharges with high plasma currents ($I_p \geq 2.0$ MeV) and monotonic q-profiles. In low plasma current discharges and in the discharges with large on-axis current holes (and, hence, extreme central magnetic shear reversal) the γ -ray emission decay times are shorter than the classical slowing down times, indicating an α -particle confinement degradation in such discharges in line with theoretical predictions. The obtained data provides essential information for studying α -particle confinement and for assessing the potential of the diagnostic technique in the perspective of ITER.

INTRODUCTION

The nuclear reaction $D(T,n)^4\text{He}$ between Deuterium (D) and Tritium (T) is the main source of energy in a thermonuclear fusion reactor with magnetic confinement. The power for the self-sustained DT-plasma burn is provided by the ^4He -ions (α particles) which are born with an average energy of 3.5 MeV and transfer the energy to the thermal plasma during their slowing down. Investigation of the α -particle behaviour is a crucial task for ITER [1] and the development of the magnetic fusion reactor concept. The α particles have been studied in the full-scale DT-plasma experiments on TFTR [2] and on JET [3] tokamaks, which are fusion devices, where plasma is confined in a toroidal vessel by means of an applied toroidal magnetic field and a poloidal field mainly induced by the plasma current. In these experiments several plasma diagnostics provided the measurement of temperature changes and some other effects, caused by fast α particles [4-6].

This Letter reports the first γ -ray measurements of fusion-born α particles in JET 'trace tritium' discharges, i.e. in majority deuterium plasma after seeding with a small population of tritium Neutral Beam Injection (NBI) fast ions. The γ -ray emission from the nuclear reaction $^9\text{Be}(\alpha,n\gamma)^{12}\text{C}$ is used to measure changes in the density of the fast α particles with energy $E_\alpha > 1.7$ MeV in the post-NBI period. This important diagnostic nuclear reaction has already been applied to detect the presence of the fast α -particles in JET experiments, where ion-cyclotron-resonance heating of ^4He -beam ions was used to accelerate ^4He to the MeV range [7]. In this Letter we demonstrate how a nuclear diagnostic based on the γ -ray spectrometry of the interaction between α 's and Be impurity in plasmas [8] could be used in future magnetic fusion machines to obtain essential information on the slowing down and confinement of the fast α particles.

In present JET experiments γ -ray energy spectra are measured with a calibrated bismuth germanate (BGO) scintillation detector with diameter of 75mm and a height of 75mm [8]. The detector is located in a well-shielded bunker and views the plasma quasi-tangentially. In order to reduce neutron

and γ -ray background, the front collimator is filled with polythene to a depth of 0.5m. Behind the detector there is an additional 1.5m long dump of polythene and steel. The detector line of sight lies in a horizontal plane about 30cm below the plasma magnetic axis. During these experiments the γ rays were continuously recorded with integration time 250ms over the energy range 1-28MeV, with an energy resolution of about 4% at 10MeV.

Diagnostic capabilities of the nuclear reaction ${}^9\text{Be}(\alpha, n\gamma){}^{12}\text{C}$, the significance of which for the fusion α particles has already been reported in [9], are determined by the specific reaction cross-section. The excitation functions of first two levels of the final nucleus, ${}^{12}\text{C}$, populated in this reaction, are shown in Fig.1. The resonance structures, which are clearly seen in both cases, provide the energy selectivity for the α -particle measurements. The first energy level 4.44MeV is excited by a particles with energies exceeding 1.7MeV, and the second one, 7.65MeV, is populated by α 's with energies in excess of 4MeV. The α -particle energy probability distribution for a born in a typical deuterium-plasma JET discharge due to DT- fusion reaction with a 105keV triton-beam injected ion is also presented in Fig.1. It is seen that the beam-plasma a particles can give rise to 4.44MeV gammas (4.44 \rightarrow 0 transition), and α particles in the high-energy wing of the distribution can also excite the second level, giving rise to γ -rays with energy 3.21MeV (7.65 \rightarrow 4.44 transition). As an example, Fig.2 shows two γ -ray spectra, recorded in the same discharge: the left hand side plot shows the spectrum during 300ms T-beam blip, the right one shows the spectrum just after the NBI blip. During the injection two γ -ray peaks, 4.44MeV and 3.21MeV, are observed, however, in the post-blip time-slice the 3.21MeV peak becomes rather weak. This is an effect of changes in the distribution function, i.e. the result of a shift of the high-energy tail to the low-energy range due to the α -particle slowing-down.

Clear variations in the intensity of the 4.44-MeV γ -ray emission were observed in the post-beam-blip period of many discharges. Figure 3 shows decays of the 4.44MeV γ -ray intensity, recorded by the spectrometer in discharges with different NBI heating power. The measured rate of 14MeV neutrons, which are born during the T-beam injection, is shown as well. It is important to note that the main plasma heating (deuterium NBI) is kept constant for several seconds after the T NBI-blip, ensuring steady plasma conditions. This applies to all the shots in our database. The γ -ray decays are thus measured against unchanging plasma conditions. In these experiments the duration of T-beam blips was $t_{blip} \leq 300$ ms.

Neutrons with energy that exceeds 5MeV could give rise to the background 4.44MeV γ -rays due to the nuclear inelastic scattering ${}^{12}\text{C}(n, n'\gamma){}^{12}\text{C}$. The main source of this background γ -ray emission is the polythene plug, which is placed in front of the detector and contains carbon as main chemical element in the compound. Extraction of the neutron background is an important part of the data processing. This factor is the main source of uncertainties in the interpretation of the present measurements.

More than 20 discharges were analysed, comparing two parameters: t_g - decay time of the 4.44MeV γ -ray intensity from the reaction ${}^9\text{Be}(\alpha, n\gamma){}^{12}\text{C}$, and the characteristic classical slowing-down time of the fast α particles on electrons, t_{se} . The variation of the γ -ray intensity after the T-blip, as shown in

Fig.3, can be approximated as $I_g(t)\mu \exp(-t/t_g)$. The time variation of the 4.44MeV γ -ray emission after the end of the T-beam blip was modelled. The γ rate (R_g) is proportional to:

$$R_\gamma(t) \propto n_{Be} \int F(E_\alpha, t) \sigma(E_\alpha) v_\alpha dE_\alpha \quad (1)$$

where n_{Be} is the Be density in the plasma; $F(E_\alpha, t)$ is the slowing down energy distribution of α 's; $\sigma(E_\alpha)$ is the energy dependence of the reaction cross-section for ${}^9\text{Be}(\alpha, n\gamma){}^{12}\text{C}$; v_α is α -particle velocity. The slowing down distribution in equation (1) was assumed to follow the classical formula with slowing down on the plasma electrons characterised by time $\tau_{se} \propto T_e^{3/2}/n_e$ [10], where T_e and n_e are the temperature and density of electrons in plasma. The distribution function $F(E_\alpha, t)$ would of course be altered by any non-classical α -losses. In the case $\tau_{se} < t_{blip}$, modelling revealed that the expected rate $R_\gamma(t)$ was approximately exponential with time constant $\sim \tau_{se}$. This follows from the fact that the expected distribution function $F(E_\alpha, t)$ is a result of the α -particle slowing down within the T-beam blip period close to the function related to the case $\tau_{se} \ll t_{blip}$ shown in Fig.1. In the case $\tau_{se} \gg t_{blip}$, the initial α -distribution after the blip and prior to the measurement phase is similar to the α -particle source distribution, which is presented in Fig.1. A simple exponential decay for $R_\gamma(t)$ would not then be observed, but these conditions did not apply in any of the experimental shots studied.

Results of the comparison of measured τ_γ against calculated classical τ_{se} for the plasmas are presented in Fig.4. For the calculation of the slowing-down time, τ_{se} , the electron density $n_e(0)$ and the electron temperature $T_e(0)$, measured in the plasma centre were used. One can see from the figure that in most of the discharges with toroidal magnetic field and plasma current in the ranges 2.25-3.2T and 2.0-3.0MA the MeV α -particle slowing-down is characterised by the scaling $\tau_\gamma \approx \tau_{se}$. The discharges displaying classical behaviour are all ELMy H-modes with monotonic q profiles. Modelling of the fast α -particle slowing-down was performed using the TRANSP code [11] for several discharges. The results of these calculations are in agreement with the experimental data, within the error bars.

There are two groups of discharges, which do not follow the classical behaviour, i.e. have $\tau_\gamma \ll \tau_{se}$. One of the explanations of the fast decay of the γ -ray emission in these plasmas is the effect of a poor α -particle confinement due to the significant orbit losses. Our modelling assessments show that a minimal critical plasma current $I_{cr} > 1.5\text{-}2\text{MA}$ is required to avoid significant First Orbit (FO) losses of 3.5MeV alphas in the discharges with monotonic plasma currents. Therefore, in discharges with $I_p^{max} = 1\text{MA}$ the γ -ray decay-times are expected to be lower than τ_{se} . Another similar anomalous behaviour of the γ -ray emission decay was observed in discharges with hollow current profiles. These discharges with Internal Transport Barriers (ITB) have strongly reversed magnetic shear in the plasma centre. Measurements based on the Motional Stark Effect (MSE) [12], show very small central current density in the plasma core area, the so called 'current hole', in these discharges. The typical size of the current holes in the analysed discharges is around $0.35a$, where a is a minor radius of the plasma. According to the confinement criteria [13], developed for a particles born in a plasma with a hollow current profile, the current hole effect is equivalent to an increase of the critical plasma current, I_{cr}

$\approx 1.5/(1-x_h^{1/2})(\text{MA})$, where $x_h=r_h/a$ is an effective radial size of the current hole. For the discharges with the current hole as large as $x_h \approx 0.35$ the critical current value is equal to $I_{cr} \approx 3.7\text{MA}$. In the case of discharges with $I_p^{max} = 2 - 2.5\text{MA}$ the FO losses of 3.5MeV alphas are therefore rather significant, 20-30%. These estimates explain the faster observed decay-times of the γ -ray emission measured in the discharges with calculated slowing time for the fast α particles derived classically.

To summarise, the time-dependent γ -ray spectra from the nuclear reaction between the fast fusion α -particles and Be-impurities, ${}^9\text{Be}(\alpha, \text{ng}){}^{12}\text{C}$, were measured for the first time in deuterium-tritium plasmas. The time evolution of the MeV α -particle density was obtained, and a correlation between the decay-time of the γ -ray emission and the characteristic a particle slowing-down time in different plasma scenarios was established. The majority of the results are consistent with classical slowing-down of the MeV α particles (with time constant τ_{se}), however in discharges with low plasma current and in discharges with hollow current profiles the α -ray emission was found to decay on a much shorter time-scale, $\tau_\gamma \ll \tau_{se}$. This is attributed to orbit losses, which determine the α -particle behaviour, and this interpretation is consistent with preliminary theoretical predictions. It is necessary to emphasise that application of this γ -ray technique with dedicated multi-channel devices could provide the time- and spatial-resolved fusion α -particle measurements in next-step fusion machines, such as ITER or other burning plasma experiments.

ACKNOWLEDGEMENTS

This work has been conducted under the European Fusion Development Agreement and was funded partly by the United Kingdom Engineering and Physical Sciences Research Council and by EURATOM. The authors wish to acknowledge fruitful discussions with A.W.Morris.

REFERENCES

- [1]. Special issue of Fusion Engineering and Design **55**, 2-3 (2001) and <http://www.iter.org/>
- [2]. R. J. Hawryluk *et al.*, Phys. Rev. Lett. **72**, 3530 (1994)
- [3]. M. Keilhacker *et al.*, Nucl. Fusion **39**, 209 (1999)
- [4]. D. S. Darrow *et al.*, Phys. Plasmas **3**, 1875 (1996)
- [5]. P. R. Thomas *et al.*, Phys. Rev. Lett. **80**, 5548 (1998)
- [6]. A. A. Korotkov *et al.*, Phys. Plasmas **7**, 957 (2000)
- [7]. M. J. Mantsinen *et al.*, Phys. Rev. Lett. **88**, 105002 (2002)
- [8]. V. G. Kiptily *et al.*, Nucl. Fusion **42**, 999 (2002)
- [9]. V. G. Kiptily, Fusion Technol. **18**, 583 (1990)
- [10]. D.L. Book., NRL Plasma Formulary, Naval Res. Lab., Washington D.C., 1990
- [11]. R. V. Budny *et al.*, Nucl. Fusion **32**, 429 (1992)
- [12]. N.C. Hawkes *et al.*, Phys. Rev. Lett. **87**, 115001 (2001)
- [13]. V. Yavorskij *et al.*, Nuclear Fusion **44**, L5-L9 (2004)




Figure 1: Excitation functions of ^{12}C levels, 4.44 MeV and 7.65 MeV, which are populated in the reaction $^9\text{Be}(\alpha, \gamma)^{12}\text{C}$. The α -particle source energy distribution was calculated for the 105 keV tritium beam injected in 6 keV deuterium plasma. The steady-state α -particle energy distribution calculated for the case $\tau_{\text{se}} \ll \tau_{\text{blip}}$. Both distribution functions have arbitrary normalisation.




Figure 2: Gamma-ray spectra measured in 2.0 MA/2.25 T Pulse No: 61046 with deuterium 15 MW NBI heating and tritium 300 ms blip $P_{\text{TNBI}} \approx 1.5 \text{ MW}$; $T_e(0) \approx 6 \text{ keV}$, $n_e(0) \approx 6 \cdot 10^{19} \text{ m}^{-3}$.




Figure 3: Comparison of time evolutions of 4.44MeV γ -ray emission measured in the Pulse No's: 61044 and 61048. Pulse No: 61044: 2.0MA/2.25T, $P_{\text{DNBI}} \approx 14.5\text{MW}$ $P_{\text{TNBI}} \approx 1.5\text{MW}$; $T_e(0) \approx 5\text{keV}$, $n_e(0) \approx 4.8 \cdot 10^{19}\text{m}^{-3}$. Pulse No. 61048: 2.0MA/2.25T, $P_{\text{DNBI}} \approx 2.9\text{MW}$ $P_{\text{TNBI}} \approx 2.3\text{MW}$; $T_e(0) \approx 3.5\text{keV}$, $n_e(0) \approx 3.2 \cdot 10^{19}\text{m}^{-3}$.




Figure 4: Measured 4.44MeV γ -ray decay-times, τ_γ , for different plasma scenarios: $B_T=2.25\text{--}3.2\text{T}$, $I_p=1.0\text{--}3.0\text{MA}$. A characteristic time of evolution of the α -particle energy distribution ($E_\alpha > 1.7\text{MeV}$) calculated by TRANSP is shown for two discharges. The slowing-down time, τ_{se} , was calculated with the electron density $n_e(0)$ and the electron temperature $T_e(0)$, measured in the plasma centre.